

A GENERAL HSU-ROBBINS-ERDŐS TYPE ESTIMATE OF TAIL PROBABILITIES OF SUMS OF INDEPENDENT IDENTICALLY DISTRIBUTED RANDOM VARIABLES

ALEXANDER R. PRUSS

ABSTRACT. Let $X_1, X_2, \dots$ be a sequence of independent and identically distributed random variables, and put $S_n = X_1 + \dots + X_n$. Under some conditions on the positive sequence τ_n and the positive increasing sequence a_n , we give necessary and sufficient conditions for the convergence of $\sum_{n=1}^{\infty} \tau_n P(|S_n| \geq \varepsilon a_n)$ for all $\varepsilon > 0$, generalizing Baum and Katz's (1965) generalization of the Hsu-Robbins-Erdős (1947, 1949) law of large numbers, also allowing us to characterize the convergence of the above series in the case where $\tau_n = n^{-1}$ and $a_n = (n \log n)^{1/2}$ for $n \geq 2$, thereby answering a question of Spătaru. Moreover, some results for non-identically distributed independent random variables are obtained by a recent comparison inequality. Our basic method is to use a central limit theorem estimate of Nagaev (1965) combined with the Hoffman-Jørgensen inequality (1974).

1. INTRODUCTION AND MAIN RESULT

Hsu and Robbins [8] showed that if $X_1, X_2, \dots$ are independent and identically distributed mean zero random variables with finite variance, then

$$(1) \quad \sum_{n=1}^{\infty} P(|S_n| \geq \varepsilon n) < \infty, \quad \forall \varepsilon > 0,$$

Date: April 1, 2002.

1991 *Mathematics Subject Classification.* 60F05, 60F10, 60F15. Secondary 60E15.

Key words and phrases. Rates of convergence in the law of large numbers, complete convergence, weak mean domination, Hsu-Robbins-Erdős law of large numbers, sums of independent random variables, tail probabilities.

where $S_n = X_1 + \cdots + X_n$. Erdős [3, 4] then showed that a converse implication also holds, namely that if (1) holds and the $\{X_i\}$ are independent and identically distributed, then $E[X_1^2] < \infty$ and $E[X_1] = 0$. Since then, a number of extensions in several directions have been proved; see [13] and [17] for partial bibliographies and brief discussions.

The purpose of the present paper is to prove a new and very general extension of the Hsu-Robbins-Erdős law of large numbers. Among other results, this extension will allow one to give necessary and sufficient conditions (see Corollary 4 in Section 2) for the truth of

$$\sum_{n=1}^{\infty} n^{-1} P(|S_n| \geq \varepsilon(n \log n)^{1/2}) < \infty, \quad \forall \varepsilon > 0,$$

where $S_n = X_1 + \cdots + X_n$ for independent and identically distributed $X_1, X_2, \dots$, thereby answering a question of Spătaru.

Throughout, terms like “positive” and “increasing” indicate the non-strict varieties (“non-negative” and “non-decreasing”, respectively).

To state our general result, we need a definition: A positive sequence $\{\tau_n\}_{n=1}^{\infty}$ is said to *satisfy Condition A* provided that for all positive decreasing sequences $\{c_n\}_{n=1}^{\infty}$ such that $\sum_{n=1}^{\infty} \tau_n \min(nc_n, 1)$ converges, we likewise have $\sum_{n=1}^{\infty} \tau_n nc_n$ converging. The following Proposition gives us a simple sufficient criterion for Condition A. A proof will be given in Section 3, below.

Proposition 1. *Suppose that either $\liminf_{n \rightarrow \infty} \tau_n > 0$ or that both of the following two conditions are satisfied:*

- (a) $\liminf_{n \rightarrow \infty} n\tau_n > 0$, and
- (b) *there is a constant $C \in (0, \infty)$ such that for each $n \in \mathbb{Z}_0^+$, if $2^{n-1} \leq k < 2^n$, then*

$$C\tau_{2^{n-1}} \geq \tau_k \geq C^{-1}\tau_{2^n}.$$

Then, $\{\tau_n\}$ satisfies Condition A.

We now state our main result. Conditions (2) and (3) in the result below are technical conditions that will be satisfied for a wide class of sequences $\{\tau_n\}$ and $\{a_n\}$. In particular, they will automatically hold in the case where $a_n = L(n)n^\alpha$ and $\tau_n = K(n)n^\beta$, assuming L and K are slowly-varying functions, $\alpha > 1/3$ and $\liminf n\tau_n > 0$ (see Proposition 2, below).

Theorem 1. *Let $X_1, X_2, \dots$ be independent identically distributed random variables. Put $S_n = X_1 + \dots + X_n$. Let $\{\tau_n\}_{n=1}^\infty$ be any positive sequence of numbers, and let $\{a_n\}_{n=1}^\infty$ be an increasing strictly positive sequence tending to ∞ . Furthermore, assume there exist finite constants $\theta \geq 1$, $C > 0$ and $N \geq 2$ such that for all $n \geq N$ we have:*

$$(2) \quad \frac{a_n^{3\theta}}{n^{\theta-1}} \sum_{k=n}^{\infty} \frac{k^\theta \tau_k}{a_k^{3\theta}} \leq C \sum_{k=1}^{n-1} k \tau_k.$$

If $\theta > 1$, then additionally assume that

$$(3) \quad \liminf_{n \rightarrow \infty} \inf_{k \geq n} \frac{a_k^3}{k a_n^3} \sum_{j=1}^{n-1} j \tau_j > 0.$$

Then, if all of the following three conditions hold:

- (i) there is a sequence $\{\mu_n\}_{n=1}^\infty$ with μ_n a median of S_n for all n , such that for all $\varepsilon > 0$, we have $\sum_{n \in M(\varepsilon)} \tau_n < \infty$, where $M(\varepsilon) = \{n \in \mathbb{Z}^+ : |\mu_n| > \varepsilon a_n\}$,
- (ii) $\sum_{n=1}^\infty n \tau_n P(|X_1| \geq \varepsilon a_n) < \infty$ for all $\varepsilon > 0$, and
- (iii) $\sum_{n=1}^\infty \tau_n e^{-\varepsilon^2 a_n^2 / (n T_{\varepsilon,n})} < \infty$ for all $\varepsilon > 0$, where $T_{\varepsilon,n} = E[X_1^2 \cdot 1_{\{|X_1| < \varepsilon a_n\}}]$, and where $e^{-t/0} = 0$ for all $t > 0$,

we will have:

$$(4) \quad \sum_{n=1}^\infty \tau_n P(|S_n| \geq \varepsilon a_n) < \infty, \quad \forall \varepsilon > 0,$$

and, conversely, if the sequence $\{\tau_n\}$ satisfies Condition A and (4) holds, conditions (i), (ii) and (iii) will also hold.

The proof will be given in Section 3, below. The result is closely related to work of Klesov [12, Theorem 4], though we are working with a more general class of sequences $\{a_n\}$. Our proof will be based upon Klesov's Hoffman-Jørgensen inequality based approach, combined with a central limit theorem estimate of Nagaev [16], the latter being used rather like in [18].

The particular newness of the result is that it works for a_n near the critical growth $n^{1/2}$ involved in the central limit theorem. For

instance, as already advertised, in Corollary 4 we will use the Theorem to characterize the cases where (4) holds with $\tau_n = n^{-1}$ and $a_n = (n \log n)^{1/2}$ (for $n \geq 2$).

Now, recall that a measurable function ϕ on $[0, \infty)$ is slowly varying (in the sense of Karamata) providing that for all $\lambda > 0$ we have $\lim_{x \rightarrow \infty} \phi(\lambda x)/\phi(x) = 1$.

Proposition 2. *If K and L are strictly positive slowly varying functions on $[0, \infty)$, and if $a_n = K(n)n^\alpha$ for some $\alpha > \frac{1}{3}$ and $\tau_n = L(n)n^\beta$ for some real β such that $\liminf_n n\tau_n > 0$, then conditions (2) and (3) will be satisfied for a sufficiently large θ .*

Sketch of proof of Proposition 2. First, for sufficiently large θ , approximate $\sum_{k=n}^{\infty} a_k^{-3\theta} k^\theta \tau_k$ by an integral and use the fact that if ϕ is slowly varying then $\int_x^\infty t^{-p} \phi(t) dt$ is asymptotic to $(p-1)^{-1} x^{1-p} \phi(x)$ for large x if $p > 1$ by Karamata's integral theorem [2, Proposition 1.5.10]. It is here that the condition that $\alpha > \frac{1}{3}$ will be used; one will also need to use the easy fact that powers and products of slowly varying functions are slowly varying. This will show via a straightforward computation that the left hand side of (2) is asymptotic to $Cn^{2+\beta}K(n)$. But also using an integral to approximate the right hand side of (2) and using the fact that if ϕ is slowly varying then $\int_1^x t^p \phi(t) dt$ is asymptotic to $(p+1)^{-1} x^{p+1} \phi(x)$ for large x by Karamata's other integral theorem [2,

Proposition 1.5.11] if $p > -1$, we can see that the right hand side of (2) is also asymptotic to $Cn^{2+\beta}K(n)$, and so (2) holds.

Condition (3) can be obtained by noting that K can without loss of generality be replaced by a normalized slowly varying function (see [2, p. 15]) asymptotic to K , and then a_k^3/k will be an increasing function for sufficiently large k if $\alpha > 1/3$, by the Bojanic-Karamata theorem [2, Theorem 1.5.5], so that the infimum on the left hand side of (3) will be attained at $k = n$, and the truth of (3) will follow from the condition that $\liminf n\tau_n > 0$. Condition A will also be satisfied by $\{\tau_n\}$ under the above circumstances, as is very easy to see by Proposition 1. $\square$

Remark 1. By a maximal inequality of Montgomery-Smith [15], condition (4) is equivalent to:

$$\sum_{n=1}^{\infty} \tau_n P\left(\left|\sup_{1 \leq k \leq n} S_k\right| \geq \varepsilon a_n\right) < \infty, \quad \forall \varepsilon > 0.$$

Remark 2. Readers familiar with Hsu-Robbins-Erdős laws of large numbers may be surprised at condition (iii) in Theorem 1, since normally these laws of large numbers simply have (4) equivalent to (i) and (ii) (under appropriate conditions on $\{\tau_n\}$ and $\{a_n\}$). In general, condition (iii) cannot be eliminated from Theorem 1, and seems to become particularly significant for a_n near the critical growth $n^{1/2}$. (Example 1 in Section 2 will show that condition (iii) cannot be eliminated if $\tau_n = n^{-1}$ and $a_n = (n \log n)^{1/2}$.) However, in a number of

special cases, condition (iii) can indeed be removed, as is seen in the following result.

Theorem 2. *Suppose the $\{\tau_n\}$ is positive and $\{a_n\}$ is increasing and strictly positive, and that there are finite real constants $\theta \geq 1$, $C > 0$ and $N \geq 2$ such that for all $n \geq N$ we have:*

$$(5) \quad \frac{a_n^{2\theta}}{n^{\theta-1}} \sum_{k=n}^{\infty} \frac{k^\theta \tau_k}{a_k^{2\theta}} \leq C \sum_{k=1}^{n-1} k \tau_k$$

and

$$(6) \quad \liminf_{n \rightarrow \infty} \inf_{k \geq n} \frac{a_k^2}{k a_n^2} \sum_{j=1}^{n-1} j \tau_j > 0.$$

Then, any random variable X_1 satisfying condition (ii) of Theorem 1 automatically satisfies condition (iii) of that Theorem.

The proof will be given in Section 3, below. Note that (5) automatically implies (2), and (6) likewise implies (3).

Remark 3. It is not difficult to see that if K and L are slowly varying strictly positive functions on $[0, \infty)$, and if $a_n = K(n)n^\alpha$ for some $\alpha > \frac{1}{2}$ while $\tau_n = L(n)n^\beta$ for some real β such that $\liminf_n n\tau_n > 0$, then conditions (5) and (6) will be satisfied for a sufficiently large θ . Since by Proposition 2, conditions (2) and (3) would allow α to be $\frac{1}{2}$ (or in fact any value greater than $\frac{1}{3}$), this helps to further illustrate how condition (iii) becomes relevant close to the critical growth $n^{1/2}$ of a_n ,

but for faster growths (i.e., around n^α for $\alpha > \frac{1}{2}$), it can be eliminated by Theorem 2.

2. COROLLARIES AND APPLICATIONS

We now obtain the following generalization due to Baum and Katz [1] of the Hsu-Robbins-Erdős result, thereby showing that Theorem 1 is indeed more general than the Hsu-Robbins-Erdős law of large numbers. (See [13] for a discussion of the pedigree of the Baum and Katz result.)

Corollary 1. *Suppose that $X_1, X_2, \dots$ are independent identically distributed random variables. Let $S_n = X_1 + \dots + X_n$. Fix $r \geq 1$ and $0 < p < 2$. Then the conjunction of the conditions*

(a) *if $p \geq 1$ then $E[X_1] = 0$, and*

(b) *$E[|X_1|^p] < \infty$*

holds if and only if

$$(7) \quad \sum_{n=1}^{\infty} n^{r-2} P(|S_n| \geq \varepsilon n^{1/p}) < \infty, \quad \forall \varepsilon > 0.$$

Proof. Let $\tau_n = n^{r-2}$ and $a_n = n^{1/p}$. Observe that (2), (3), (5), (6) and Condition A all hold for appropriate choices of θ (one can use Proposition 2 and Remark 3 here if one so desires, but in fact a direct verification is easy). Then by the Marcinkiewicz-Zygmund strong law of large numbers [14, §16.4.A.3], if (a) and (b) hold, then $S_n/a_n \rightarrow 0$ almost surely, hence also in probability, and therefore condition (i) of

Theorem 1 holds as well. Moreover, (ii) is equivalent to (b), and (ii) implies (iii) by Theorem 2. Thus, by Theorem 1, if (a) and (b) hold, (7) follows. Conversely, if (7) holds, then since τ_n satisfies Condition A by Proposition 1, it follows from Theorem 1 that (i) and (ii) (and (iii), but that is not needed) hold. Hence (b) holds, since it is equivalent to (ii). It remains to show that (a) holds. The easiest way to do this is to note that if $p \geq 1$ then by the Marcinkiewicz-Zygmund strong law of large numbers, we have $(S_n - nE[X_1])/a_n \rightarrow 0$ almost surely, hence in probability, and therefore $(\mu_n - nE[X_1])/n^{1/p} \rightarrow 0$, where μ_n is a median of S_n , and by condition (i) it will then follow that $E[X_1] = 0$. $\square$

Recall that random variables $X_1, \dots, X_n$ are said to be *K-weakly mean dominated* by a random variable X providing that for all λ we have:

$$\frac{1}{n} \sum_{k=1}^n P(|X_k| \geq \lambda) \leq KP(|X| \geq \lambda)$$

(see [6]).

Corollary 2. *Let X be a random variable and K any finite constant.*

Let $\{X_{k,n}\}_{1 \leq k \leq n; n \geq 1}$ be a triangular array of random variables such that $X_{1,n}, \dots, X_{n,n}$ are independent and K -weakly mean dominated by X for each fixed n . Let $\{\tau_n\}$ be a positive sequence of numbers and let $\{a_n\}$ be an increasing strictly positive sequence for $n \geq 1$. Put

$S_n = X_{1,n} + \cdots + X_{n,n}$. Suppose that condition (i) of Theorem 1 holds, (2) is satisfied and if $\theta > 1$ then (3) holds as well. Assume that conditions (ii) and (iii) of Theorem 1 are satisfied with X in place of X_1 . Then,

$$\sum_{n=1}^{\infty} \tau_n P(|S_n| \geq \varepsilon a_n) < \infty$$

for all $\varepsilon > 0$.

Proof. By Theorem 1 we have

$$(8) \quad \sum_{n=1}^{\infty} \tau_n P(|S'_n| \geq \varepsilon a_n) < \infty, \quad \forall \varepsilon > 0,$$

where S'_n is the sum of n independent copies of X (condition (i) in this case will hold trivially by symmetry, while (ii) and (iii) were assumed in the statement of Corollary 2.) One may slightly modify the comparison result in [20, Corollary 1] by assuming our Theorem 1's condition (i) in place of the assumption in that paper that $S_n/a_n \rightarrow 0$ in probability, which modification only very slightly affects the proof (one will need to use (38), below, after obtaining the convergence of [20, series (1.2)] in the original proof in [20]). Thus modified, [20, Corollary 1] together with (8) yields the conclusion of our Corollary 2. $\square$

The following Corollary yielding a result similar to one of Hu, Moricz and Taylor [9] (cf. [6, 10, 11]) can be derived from Corollary 2 exactly in the way that Corollary 1 was derived from Theorem 1.

Corollary 3. *Let X be a random variable and let K be any finite constant. Let $\{X_{k,n}\}_{1 \leq k \leq n; n \geq 1}$ be a triangular array of random variables such that $X_{1,n}, \dots, X_{n,n}$ are independent K -weakly mean dominated by X . Let $S_n = X_{1,n} + \dots + X_{n,n}$. Fix $r \geq 1$ and $0 < p < 2$. Suppose $E[|X|^{rp}] < \infty$. If $p \geq 1$ then assume also that $E[X_{n,1} + \dots + X_{n,n}] = 0$ for all n . Then,*

$$\sum_{n=1}^{\infty} n^{r-2} P(|S_n| \geq \varepsilon n^{1/p}), \quad \forall \varepsilon > 0.$$

Now define $\log^+ x = \log(2 + x)$. It is easy to see that Theorem 1 implies the following result.

Corollary 4. *Let $X_1, X_2, \dots$ be independent and identically distributed random variables. Then,*

$$(9) \quad \sum_{n=2}^{\infty} n^{-1} P(|S_n| \geq \varepsilon (n \log n)^{1/2}) < \infty, \quad \forall \varepsilon > 0,$$

if and only if all of the following three conditions hold:

- (a) $E[X_1] = 0$,
- (b) $E[X_1^2 / \log^+ |X_1|] < \infty$, and
- (c) $\sum_{n=2}^{\infty} n^{-1-\varepsilon^2/T_{\varepsilon,n}} < \infty$ for all $\varepsilon > 0$, where $T_{\varepsilon,n} = E[X_1^2 \cdot 1_{\{|X_1| < \varepsilon (n \log n)^{1/2}\}}]$.

The proof of the following Lemma will be given at the end of Section 3.

Lemma 1. *If condition (b) of Corollary 4 holds, then $(S_n - E[S_n])/(n \log n)^{1/2} \rightarrow 0$ in probability as $n \rightarrow \infty$.*

Remark 4. It is not known whether “in probability” can be replaced by “almost surely” in Lemma 1.

Proof of Corollary 4. Let $\tau_n = 1/n$ and $a_n = (n \log n)^{1/2}$ for $n \geq 2$. Note that (2) is easily seen to be satisfied with $\theta = 1$, and Condition A holds by Proposition 1. Observe that (b) is equivalent to condition (ii) of Theorem 1 in the present setting, and that if (b) holds, then (a) is equivalent to (i) by Lemma 1, so that the conjunction of conditions (a) and (b) of Corollary 4 is equivalent to that of conditions (i) and (ii) of Theorem 1. Also, it is easy to check that (c) is equivalent to (iii), since the summands in the sums in both conditions are equal. $\square$

Professor Aurel Spătaru has asked the author whether (9) is equivalent to the conjunction of (a) and (b). This would be expected by analogy with other Hsu-Robbins-Erdős laws of large numbers (such as Corollary 1). It is this question that has inspired the present paper. In light of the Corollary 4, Spătaru’s question is equivalent to asking whether the conjunction of (a) and (b) implies (c). The following counterexample that Professor Stephen Montgomery-Smith has privately communicated to the author shows that the answer is negative, and hence so is the answer to Spătaru’s question.

Example 1. Let $\{K_m\}_{m=0}^\infty$ be a very rapidly increasing strictly positive sequence, with $K_0 = 0$. The degree of rapidity of increase will be chosen later so as to make the argument go. Let $\psi(t) = (t \log t)^{1/2}$ for $t \geq 2$. Extend ψ linearly to the interval $[0, 2]$ in such a way that $\psi(0) = 0$. Let ϕ be the inverse function of ψ . Assume that $K_1 \geq 2$. Let X_1 be a random variable such that $P(X_1 = \psi(K_m)) = P(X_1 = -\psi(K_m)) = 2^{-m-1}/K_m$ for all m , and with $P(|X_1| \notin \{0\} \cup \{\psi(K_m) : m \in \mathbb{Z}^+\}) = 0$.

Let $\varepsilon = 1$. We have

$$E[\phi(|X_1|)] = \sum_{m=1}^{\infty} 2^{-m} K_m^{-1} \cdot K_m = 1.$$

It is easy to see that in general $E[\phi(|X_1|)] < \infty$ if and only if (b) holds, and hence indeed (b) is satisfied in the present case. So is (a), since X_1 is symmetric (i.e., X_1 and $-X_1$ have the same distribution) and so we can put $\mu_n = 0$ for all n . Now, let $M(n) = \max\{m \in \mathbb{Z}_0^+ : K_m < n\}$. Then, with $T_{1,n}$ as in condition (iii) of Theorem 1, and as $K_m \geq 2$ for all $m \geq 2$

$$\begin{aligned} T_{1,n} &= \sum_{m=1}^{M(n)} (2^{-m}/K_m) (\psi(K_m))^2 \\ &= \sum_{m=1}^{M(n)} (2^{-m}/K_m) (K_m \log K_m) \\ &= \sum_{m=1}^{M(n)} 2^{-m} \log K_m \geq 2^{-M(n)} \log K_{M(n)}. \end{aligned}$$

Note that if $K_m < n \leq K_{m+1}$, then $M(n) = m$ and so $T_{1,n} \geq$

$2^{-m} \log K_m$. Thus, for $m \geq 1$ we have

$$(10) \quad \sum_{n=K_m+1}^{K_{m+1}} n^{-1-1/T_{1,n}} \geq \sum_{n=K_m+1}^{K_{m+1}} n^{-1-2^m/\log K_m}.$$

Now, for any $K \geq 2$ and $m \in \mathbb{Z}^+$, let $L(K, m)$ be an integer greater than K and sufficiently large that:

$$(11) \quad \sum_{n=K+1}^{L(K,m)} n^{-1-2^m/\log K} \geq \frac{1}{2} \sum_{n=K+1}^{\infty} n^{-1-2^m/\log K}.$$

Such an $L(K, m)$ exists because the sum on the right hand side converges. Observe furthermore that

$$(12) \quad \sum_{n=K+1}^{\infty} n^{-1-2^m/\log K} \geq C \cdot 2^{-m} (\log K) (K+1)^{-2^m/\log K},$$

for an absolute constant $C > 0$ independent of $K \geq 2$ and $m \geq 0$.

Combining (10)–(12) we see that if $K_{m+1} \geq L(K_m, m)$, then we have

$$(13) \quad \begin{aligned} \sum_{n=K_m+1}^{K_{m+1}} n^{-1-1/T_{1,n}} &\geq \frac{C}{2} \cdot 2^{-m} (\log K_m) (K_m+1)^{-2^m/\log K_m} \\ &= 2^{-m-1} C (\log K_m) \exp \left(-2^m \frac{\log(K_m+1)}{\log K_m} \right). \end{aligned}$$

Inductively choosing the K_m so that for all m we have both

$$2^{-m-1} (\log K_m) \cdot \exp \left(-2^m \frac{\log(K_m+1)}{\log K_m} \right) \geq 1,$$

and $K_{m+1} \geq L(K_m, m)$, we then find by (13) that that

$$\sum_{n=2}^{\infty} n^{-1-1/T_{1,n}} \geq \sum_{m=1}^{\infty} \sum_{n=K_m+1}^{K_{m+1}} n^{-1-1/T_{1,n}} \geq C \sum_{m=1}^{\infty} 1 = \infty.$$

Hence Corollary 4's condition (c) fails, and so we do have our desired counterexample satisfying (a), (b) but not (c), and hence by that Corollary, with inequality (9) also failing.

Although the answer to Spătaru's question is negative, we do have the result under a slightly stronger moment condition than Corollary 4's condition (b).

Corollary 5. *Let X be a random variable and let K be any finite constant. Let $\{X_{k,n}\}_{1 \leq k \leq n; n \geq 1}$ be a triangular array of random variables such that $X_{1,n}, \dots, X_{n,n}$ are K -weakly mean dominated by X . Assume that*

- (a) $E[X_{n,1} + \dots + X_{n,n}] = 0$ for all n , and
- (b) $E \left[\frac{X^2 (\log^+ \log^+ |X|)^{1+\delta}}{\log^+ |X|} \right] < \infty$ for some $\delta > 0$.

Then,

$$\sum_{n=2}^{\infty} \frac{1}{n} P(|S_n| \geq \varepsilon (n \log n)^{1/2}) < \infty, \quad \forall \varepsilon > 0.$$

Proof. Using Corollary 2 and the same methods as in Corollary 4 it suffices to show that if condition (b) of Corollary 5 holds, then condition (c) of Corollary 4 is satisfied with X in place of X_1 . To show this, without loss of generality (rescaling X if necessary) assume $\varepsilon = 1$.

Note that if $T_{1,n} = E[X^2 \cdot 1_{\{|X| < a_n\}}]$, where $a_n = (n \log n)^{1/2}$, then

$$\begin{aligned} T_{1,n} &\leq E \left[\frac{X^2 (\log^+ \log^+ |X|)^{1+\delta}}{\log^+ |X|} \right] \cdot \frac{\log^+ a_n}{(\log^+ \log^+ a_n)^{1+\delta}} \\ (14) \quad &\leq C \frac{\log^+ a_n}{(\log^+ \log^+ a_n)^{1+\delta}} \\ &\leq C' \frac{\log^+ n}{(\log^+ \log^+ n)^{1+\delta}}, \end{aligned}$$

where δ is as in (b), while C and C' are strictly positive finite constants independent of $n \geq 2$ (but dependent on the value of the expectation

in (b)). Hence, if $N \geq 2$ is sufficiently large that $(\log^+ \log^+ n)^{1+\delta} \geq$

$2C' \log \log n$ for all $n \geq N$, then we have:

$$\begin{aligned} \sum_{n=N}^{\infty} n^{-1-1/T_{1,n}} &= \sum_{n=N}^{\infty} n^{-1} \exp(-(\log n)/T_{1,n}) \\ &\leq \sum_{n=N}^{\infty} n^{-1} \exp(-(\log^+ \log^+ n)^{1+\delta}/C') \\ &\leq \sum_{n=N}^{\infty} n^{-1} \exp(-2 \log \log n) = \sum_{n=N}^{\infty} \frac{1}{n(\log n)^2} < \infty, \end{aligned}$$

by (14) and the choice of N . Thus condition (c) of Corollary 4 is indeed satisfied with X in place of X_1 . $\square$

3. PROOFS AND AUXILIARY RESULTS

Proof of Proposition 1. If $\liminf \tau_n > 0$ and $\sum_{n=1}^{\infty} \tau_n \min(1, nc_n) < \infty$, then there are only finitely many $n \in \mathbb{Z}^+$ for which $nc_n \geq 1$, and hence $\sum_{n=1}^{\infty} \tau_n nc_n$ must also converge since it differs from $\sum_{n=1}^{\infty} \tau_n \min(1, nc_n)$ only in finitely many terms.

It remains to show that if (a) and (b) hold, then Condition A. To do this, suppose c_n is a decreasing sequence such that

$$(15) \quad \sum_{n=1}^{\infty} \tau_n \min(1, nc_n) < \infty.$$

Then, using (b) we have:

$$\begin{aligned} (16) \quad \infty &> \sum_{n=1}^{\infty} \tau_n \min(1, nc_n) \\ &\geq \sum_{k=0}^{\infty} C^{-1} 2^k \tau_{2^{k+1}} \min(1, 2^k c_{2^{k+1}}) \\ &\geq \frac{1}{2C} \sum_{k=0}^{\infty} 2^k \tau_{2^{k+1}} \min(1, 2^{k+1} c_{2^{k+1}}). \end{aligned}$$

Now, $\liminf 2^k \tau_{2^{k+1}} > 0$ by (a), and hence it follows that only finitely many of the $\min(1, 2^{k+1} c_{2^{k+1}})$ can equal 1 (since otherwise the right hand side of (16) would be infinite), so that except for at most finitely many values of k , we have $\min(1, 2^{k+1} c_{2^{k+1}}) = 2^{k+1} c_{2^{k+1}}$. It thus follows from (16) and (b) that

$$\begin{aligned} \infty &> \frac{1}{2} \sum_{k=0}^{\infty} 2^{k+1} \tau_{2^{k+1}} \cdot 2^{k+1} c_{2^{k+1}} \\ &\geq \frac{1}{2C} \sum_{k=0}^{\infty} \sum_{j=0}^{2^{k+1}-1} \tau_{j+2^{k+1}} \frac{j+2^{k+1}}{2} \cdot c_{j+2^{k+1}} \\ &= \frac{1}{4C} \sum_{n=2}^{\infty} \tau_n n c_n, \end{aligned}$$

and the proof is complete. $\square$

Lemma 2. *In the setting of Theorem 1, if the series $\{\tau_n\}$ satisfies Condition A, then condition (4) entails (i) and (ii).*

Proof. Assume (4) holds. Fix $\varepsilon > 0$ and any sequence of medians μ_n of the S_n . Let M be as in condition (i). Then, it is easy to see that for $n \in M$ we have $P(|S_n| \geq \varepsilon a_n) \geq 1/2$. The convergence of $\sum_{n \in M} \tau_n$ follows immediately from this and (4), and so condition (i) holds.

On the other hand, by (4) and the remark following the main theorem in [19], we have

$$\sum_{n=1}^{\infty} \tau_n \min(1, nP(|X_1| \geq \varepsilon a_n)) < \infty$$

for all $\varepsilon > 0$. Condition (ii) follows immediately from this together with the fact that $\{\tau_n\}$ satisfies Condition A while $\{a_n\}$ is increasing. $\square$

Lemma 3. *Suppose that $\{\tau_n\}$ and $\{\rho_n\}$ are positive and that $\{b_n\}$ is increasing and strictly positive. Let X be any random variable. If there is a constant $C \in (0, \infty)$ such that for all $n \geq 2$:*

$$(17) \quad b_n^t \sum_{k=n}^{\infty} \rho_k \leq C \sum_{k=1}^{n-1} k \tau_k,$$

then

$$\sum_{n=2}^{\infty} \rho_n E[|X|^t \cdot 1_{\{|X| < b_n\}}] \leq C \sum_{n=1}^{\infty} n \tau_n P(|X| \geq b_n).$$

Proof of Lemma 3. Assume without loss of generality that $b_1 = 0$.

Then, by Fubini's theorem and (17):

$$\begin{aligned} \sum_{n=2}^{\infty} \rho_n E[|X|^t \cdot 1_{\{|X| < b_n\}}] &\leq \sum_{n=2}^{\infty} \rho_n \sum_{k=1}^n b_k^t P(b_{k-1} \leq |X| < b_k) \\ &= \sum_{k=2}^{\infty} P(b_{k-1} \leq |X| < b_k) b_k^t \sum_{n=k}^{\infty} \rho_n \\ &\leq C \sum_{k=2}^{\infty} P(b_{k-1} \leq |X| < b_k) \sum_{n=1}^{k-1} n \tau_n \\ &= C \sum_{n=1}^{\infty} n \tau_n \sum_{k=n+1}^{\infty} P(b_{k-1} \leq |X| < b_k) \\ &= C \sum_{n=1}^{\infty} n \tau_n P(|X| \geq b_n). \end{aligned}$$

□

Lemma 4. *Suppose that $\{\alpha_n\}$ and $\{\beta_n\}$ are sequences in $[0, 1]$, that $\{\tau_n\}$ is a positive sequence, and that $r \in \mathbb{Z}^+$ is such that $\sum_{n=1}^{\infty} \tau_n |\alpha_n - \beta_n|^r < \infty$. If $\sum_{n=1}^{\infty} \tau_n \beta_n < \infty$, then $\sum_{n=1}^{\infty} \tau_n \alpha_n^r < \infty$.*

Proof. There is a polynomial $p_r(x, y)$ of degree $r - 1$ with coefficients depending only on r such that $(x - y)^r = x^r - y p_r(x, y)$. Let c_r be the

maximum of p_r over $[0, 1]^2$. Then:

$$\infty > \sum_{n=1}^{\infty} \tau_n |\alpha_n - \beta_n|^r = \sum_{n=1}^{\infty} \tau_n |\alpha_n^r - \beta_n p_r(\alpha_n, \beta_n)| \geq \sum_{n=1}^{\infty} \tau_n (\alpha_n^r - c_r \beta_n).$$

If $\sum_{n=1}^{\infty} \tau_n \beta_n$ converges, then it follows that $\sum_{n=1}^{\infty} \tau_n \alpha_n^r$ also converges.

□

Lemma 5. *Let $\{\tau_n\}$ be a positive sequence and let $\{b_n\}$ be a strictly positive increasing sequence for $n \geq 1$. Let X be a random variable such that*

$$(18) \quad \sum_{n=1}^{\infty} \tau_n n P(|X| \geq b_n) < \infty.$$

Fix $\nu \in [0, \infty)$. Put $T_k = \sum_{n=1}^k n \tau_n$. Suppose that there is a constant $C \in (0, \infty)$ and a $\theta \in [1, \infty)$ such that:

$$(19) \quad \frac{b_n^{\nu\theta}}{n^{\theta-1}} \sum_{k=n}^{\infty} \frac{k^\theta \tau_k}{b_k^{\nu\theta}} \leq C T_{n-1},$$

for all $n \geq 2$, and

$$(20) \quad \frac{k b_n^\nu}{b_k^\nu} \leq C T_{n-1},$$

whenever $k \geq n \geq 2$. Then:

$$\sum_{n=1}^{\infty} \tau_n \left(\frac{n E[|X|^\nu \cdot 1_{\{|X| < b_n\}}]}{b_n^\nu} \right)^\theta < \infty$$

The proof is based on methods of Klesov [12, Proof of Theorem 4].

Proof. Set $X^{(n)} = X \cdot 1_{\{|X| < b_n\}}$. Let $b_0 = 0$ and put $t_n = E[|X^{(n)}|^\nu]$ for $n \geq 0$. Note that $t_0 = 0$. Put $\delta_n = t_n^\theta - t_{n-1}^\theta$ for $n \geq 1$. For convenience, let $T_0 = 1$; redefining C if necessary, we may assume that

(19) and (20) also hold for $n = 1$. Then, since $t_n^\theta = \sum_{k=1}^n \delta_k$, and using

Fubini's theorem and (19):

$$\begin{aligned}
 \sum_{n=1}^{\infty} \tau_n \left(\frac{nE[|X^{(n)}|^\nu]}{b_n^\nu} \right)^\theta &= \sum_{n=1}^{\infty} \tau_n (nb_n^{-\nu})^\theta \sum_{k=1}^n \delta_k \\
 (21) \qquad &= \sum_{k=1}^{\infty} \delta_k \sum_{n=k}^{\infty} \tau_n (nb_n^{-\nu})^\theta \\
 &\leq C \sum_{k=1}^{\infty} \delta_k k^{\theta-1} b_k^{-\nu\theta} T_{k-1}.
 \end{aligned}$$

Let $P_k = P(b_{k-1} \leq |X| < b_k)$. Then, note that

$$(22) \qquad \delta_k \leq c(t_k - t_{k-1})t_k^{\theta-1} \leq cP_k b_k^\nu t_k^{\theta-1},$$

where c is a finite constant depending only on θ and such that $x^\theta - y^\theta \leq c(x - y)x^{\theta-1}$ whenever $0 \leq y \leq x$. Now, fix $\ell \in \mathbb{Z}^+$. Let $\rho_n = \ell/b_\ell^\nu$ for $n = \ell$ and put $\rho_n = 0$ for all other n . Observe that by (20) we have

$$(23) \qquad b_n^\nu \sum_{j=n}^{\infty} \rho_j \leq CT_{n-1},$$

for $n \leq \ell$ and the same inequality trivially holds for $n > \ell$.

By Lemma 3 (with $t = \nu$) and (23), it follows that if $\ell \geq 2$ so that

$\rho_1 = 0$, then then

$$\begin{aligned}
 \frac{lt_\ell}{b_\ell^\nu} &= \sum_{n=1}^{\infty} \rho_n t_n = \sum_{n=2}^{\infty} \rho_n E[|X^{(n)}|^\nu] \\
 &\leq C \sum_{n=1}^{\infty} n \tau_n P(|X| \geq b_n) < \infty,
 \end{aligned}$$

where the finiteness of the right hand side followed from (18). Thus,

$K \stackrel{\text{def}}{=} \sup_{\ell \geq 1} \ell t_\ell / b_\ell^\nu < \infty$. Then, $t_k \leq K b_k^\nu / k$, so that (22) yields:

$$\delta_k \leq cK^{\theta-1} P_k b_k^\nu (b_k^\nu / k)^{\theta-1} = cK^{\theta-1} \frac{P_k b_k^{\nu\theta}}{k^{\theta-1}}.$$

Putting this into (21), and recalling that $T_1 = 1$, we see that:

$$\begin{aligned}
\sum_{n=1}^{\infty} \tau_n \left(\frac{nE[|X^{(n)}|^{\nu}]}{b_n^{\nu}} \right)^{\theta} &\leq CcK^{\theta-1} \sum_{k=1}^{\infty} P_k T_{k-1} \\
&= CcK^{\theta-1} \left(P_1 + \sum_{k=2}^{\infty} P_k \sum_{n=1}^{k-1} n\tau_n \right) \\
&= CcK^{\theta-1} \left(P_1 + \sum_{n=1}^{\infty} n\tau_n \sum_{k=n+1}^{\infty} P_k \right) \\
&= CcK^{\theta-1} \left(P_1 + \sum_{n=1}^{\infty} n\tau_n P(|X| \geq b_n) \right) < \infty,
\end{aligned}$$

by Fubini's theorem and (18). $\square$

The following version of the Hoffman-Jørgensen inequality [7] will be needed for the proof of Theorem 1 in the case $\theta > 1$ and follows immediately from [13, Lemma 2.2].

Lemma 6. *Let $X_1, \dots, X_n$ be independent symmetric random variables, and let $S_n = X_1 + \dots + X_n$. Then for each $r \in \mathbb{Z}^+$ there exist finite constants C_r and D_r such that for all $\lambda \geq 0$ we have:*

$$P(|S_n| \geq \lambda) \leq C_r \sum_{k=1}^n P(|X_k| \geq \lambda/(2r)) + D_r [P(|S_n| \geq \lambda/(2r))]^r.$$

Let Φ be the distribution function of a $(0,1)$ normal random variable.

Lemma 7. *Under the global conditions of Theorem 1, suppose that condition (ii) is satisfied and that X_1 is symmetric. Then, the following four conditions are equivalent:*

- (a) $\sum_{n=1}^{\infty} \tau_n P(|S_n| \geq \varepsilon a_n) < \infty$ for all $\varepsilon > 0$;

- (b) for all $\gamma \in (0, 1]$ and $\varepsilon > 0$ we have $\sum_{n=1}^{\infty} \tau_n [1 - \Phi(\gamma \varepsilon a_n / (n T_{\varepsilon, n})^{1/2})] < \infty$, where $T_{\varepsilon, n}$ is as in Theorem 1, and where $\Phi(t/0) = 0$ for all $t > 0$;
- (c) there is an $s \geq 1$ such that for all $\varepsilon > 0$ we have $\sum_{n=1}^{\infty} \tau_n [1 - \Phi(\varepsilon a_n / (n T_{\varepsilon, n})^{1/2})]^s < \infty$;
- (d) condition (iii) of Theorem 1 holds.

The equivalence of (a) and (b) will be the most difficult part to prove, and will involve a similar method of proof to that in [18], using a central limit theorem estimate in the present case due to Nagaev [16]. In the case where $\theta > 1$, we will also need the Hoffman-Jørgensen inequality based methods of Klesov [12, Proof of Theorem 4].

Proof of Lemma 7. Note that:

$$(24) \quad 1 - \Phi(x) = \pi^{-1/2} x^{-1} e^{-x^2/2} (1 - O(x^{-2}))$$

as $x \rightarrow \infty$. Thus:

$$(25) \quad 1 - \Phi(x) \leq C e^{-x^2/2},$$

for all $x \geq 0$ and an absolute constant C .

First suppose (d) holds. Put $\gamma' = \gamma/2$. Then, (d) implies that:

$$(26) \quad \sum_{n=1}^{\infty} \tau_n e^{-(\gamma' \varepsilon)^2 a_n^2 / n T_{\gamma' \varepsilon, n}} < \infty,$$

for all $\varepsilon > 0$. Observe now that as $\gamma' < \gamma \leq 1$ and by definition of $T_{\varepsilon,n}$:

$$(27) \quad \begin{aligned} T_{\varepsilon,n} &\leq T_{\varepsilon\gamma',n} + (\varepsilon\gamma a_n)^2 P(|X_1| \geq \varepsilon\gamma' a_n) \\ &\leq \max(2T_{\varepsilon\gamma',n}, 2(\varepsilon\gamma a_n)^2 P(|X_1| \geq \varepsilon\gamma' a_n)). \end{aligned}$$

Then, by (25) and (27):

$$\begin{aligned} \sum_{n=1}^{\infty} \tau_n [1 - \Phi(\gamma\varepsilon a_n / (nT_{\varepsilon,n})^{1/2})] &\leq C \sum_{n=1}^{\infty} \tau_n e^{-\gamma^2 \varepsilon^2 a_n^2 / 2nT_{\varepsilon,n}} \\ &\leq C \max \left(\sum_{n=1}^{\infty} \tau_n e^{-\gamma^2 \varepsilon^2 a_n^2 / 4nT_{\varepsilon\gamma',n}}, \sum_{n=1}^{\infty} \tau_n e^{-1/4nP(|X_1| \geq \varepsilon\gamma' a_n)} \right), \end{aligned}$$

where we use the convention that $e^{-1/0} = 0$. Now, the first sum on the right hand side here converges by (26) since $\gamma^2/4 = (\gamma')^2$, while the second converges by the elementary inequality $e^{-1/x} \leq x$ valid for all $x \geq 0$, together with condition (ii) of Theorem 1. Hence, (d) implies (b).

The implication from (b) to (c) is trivial. Suppose now that (c) holds. Let $u_{\delta,n} = a_n / (nT_{\delta,n})^{1/2}$. Then,

$$(28) \quad \sum_{n=1}^{\infty} \tau_n e^{-s\delta^2 u_{\delta,n}^2} < \infty,$$

for all $\delta > 0$ by (c) and (24). Fix $\varepsilon > 0$ and let $\delta = \varepsilon / (2s)^{1/2}$. Then,

$u_{\delta,n} \leq u_{\varepsilon,n}$ and so

$$e^{-s\delta^2 u_{\delta,n}^2} \geq e^{-s\delta^2 u_{\varepsilon,n}^2} = e^{-\varepsilon^2 u_{\varepsilon,n}^2 / 2}.$$

Condition (d) then follows from this together with (24) and (28). Hence (c) implies (d), so that we have shown that (d) $\Rightarrow$ (b) $\Rightarrow$ (c) $\Rightarrow$ (d).

All we now need to prove is the equivalence of (a) and (b). To do this, assume we are in the setting of Theorem 1 and that (ii) holds. Fix

$\varepsilon > 0$. Let $X_{k,n}(\varepsilon) = X_k \cdot 1_{\{|X_k| < \varepsilon a_n\}}$. Let $S_n(\varepsilon) = X_{1,n}(\varepsilon) + \dots + X_{n,n}(\varepsilon)$.

Set $A_n(\varepsilon) = \bigcup_{k=1}^n \{X_k \neq X_{k,n}(\varepsilon)\}$. Observe that $S_n = S_n(\varepsilon)$ except possibly on $A_n(\varepsilon)$, and that

$$(29) \quad \sum_{n=1}^{\infty} \tau_n P(A_n(\varepsilon)) \leq \sum_{n=1}^{\infty} n \tau_n P(|X_1| \geq \varepsilon a_n) < \infty,$$

by (ii). It follows from (29) and from the equality of S_n and $S_n(\varepsilon)$ outside $A_n(\varepsilon)$ that (a) holds if and only if

$$(30) \quad \sum_{n=1}^{\infty} \tau_n P(|S_n(\varepsilon)| \geq \varepsilon a_n) < \infty, \quad \forall \varepsilon > 0.$$

We now need only show that (30) holds if and only if (b) holds, and we will be done. Note that $T_{\varepsilon,n} = E[(X_{1,n}(\varepsilon))^2]$. Fix $\gamma > 0$ to be chosen later as needed. Then, since all the $X_{k,n}(\varepsilon)$ and $S_n(\varepsilon)$ have mean zero by symmetry, and since $X_{1,n}(\varepsilon), \dots, X_{n,n}(\varepsilon)$ are identically distributed for a fixed n , by Nagaev's central limit theorem estimate [16, Theorem 3] we have:

$$\left| P(|S_n(\varepsilon)| \geq \gamma \varepsilon a_n) - 2 \left[1 - \Phi \left(\frac{\gamma \varepsilon a_n}{n T_{\varepsilon,n}^{1/2}} \right) \right] \right| \leq c n \frac{E[|X_{1,n}(\varepsilon)|^3]}{(\gamma \varepsilon a_n)^3},$$

for an absolute constant $c < \infty$. Since the left hand side never exceeds 1, it follows further that for a (possibly different) absolute constant c , we have:

$$(31) \quad \left| P(|S_n(\varepsilon)| \geq \gamma \varepsilon a_n) - 2 \left[1 - \Phi \left(\frac{\gamma \varepsilon a_n}{n T_{\varepsilon,n}^{1/2}} \right) \right] \right| \leq c \min \left(1, n \frac{E[|X_{1,n}(\varepsilon)|^3]}{(\gamma \varepsilon a_n)^3} \right),$$

Observe that if (2) holds for some θ , then it also holds for all greater values. We now have a quick proof if $\theta = 1$. For then, by (31) (with

$\gamma = 1$) we can see that the equivalence of (b) and (30) would follow as soon as we could show that we have

$$(32) \quad \sum_{n=1}^{\infty} n\tau_n \frac{E[|X_{1,n}(\varepsilon)|^3]}{a_n^3} < \infty.$$

But (32) follows from the validity of (2) for $\theta = 1$ and from condition (ii) of Theorem 1, by an application of Lemma 3 with $\rho_n = n\tau_n/a_n^3$, $X \equiv X_1$, $t = 3$ and $b_n = \varepsilon a_n$.

Suppose now we are working with $\theta > 1$, so that (3) also holds. Let r be an integer greater than or equal to θ . I now claim that:

$$(33) \quad \sum_{n=1}^{\infty} \tau_n \left(\min \left(1, \frac{nE[|X_{1,n}(\varepsilon)|^3]}{a_n^3} \right) \right)^r < \infty, \quad \forall \varepsilon > 0.$$

Suppose for now that this has been shown. If (a) holds, then as noted before, (30) does likewise. Letting $\alpha_n = 2[1 - \Phi(\varepsilon a_n/(nT_{\varepsilon,n})^{1/2})]$ and $\beta_n = P(|S_n(\varepsilon)| \geq \varepsilon a_n)$, we see that by (30) together with (33) and (31) (with $\gamma = 1$), we do have the conditions of Lemma 4 satisfied, so that $\sum_{n=1}^{\infty} \tau_n \alpha_n^r < \infty$, and hence (c) follows, whence (b) follows by the already proved equivalence of (b), (c) and (d).

Conversely, suppose (b) holds. This time letting $\beta_n = 2[1 - \Phi(\gamma \varepsilon a_n/(nT_{\varepsilon,n})^{1/2})]$ and $\alpha_n = P(|S_n(\varepsilon)| \geq \gamma \varepsilon a_n)$, using (b), together with (31), (33) and Lemma 4, we see that $\sum_{n=1}^{\infty} \tau_n \alpha_n^r < \infty$, i.e.,

$$(34) \quad \sum_{n=1}^{\infty} \tau_n [P(|S_n(\varepsilon)| \geq \gamma \varepsilon a_n)]^r < \infty.$$

Let $\gamma = (2r)^{-1}$. By the Hoffman-Jørgensen inequality (Lemma 6), we have:

$$\begin{aligned} \sum_{n=1}^{\infty} \tau_n P(|S_n(\varepsilon)| \geq \varepsilon a_n) \\ \leq C_r \sum_{n=1}^{\infty} \tau_n P(|X_n(\varepsilon)| \geq \gamma \varepsilon a_n) + D_r \sum_{n=1}^{\infty} \tau_n [P(|S_n(\varepsilon)| \geq \gamma \varepsilon a_n)]^r. \end{aligned}$$

It is easy to see that the first sum on the right hand side is no greater than $\sum_{n=1}^{\infty} \tau_n P(A_n(\gamma\varepsilon))$, which converges by (29), and the second converges by (34). Hence, (30) follows, and as already shown this implies (a).

Hence, all we need to show is that (33) holds. Changing finitely many values of a_n and τ_n , we may assume that (2) holds for all $n \geq 2$, and that (3) gives:

$$(35) \quad \sup_{k \geq n} \frac{ka_n^3}{a_k^3} \leq C' \sum_{k=1}^{n-1} n\tau_k,$$

for all $n \geq 2$ and a finite C' . We can now apply Lemma 5 with $\nu = 3$, $b_n = \varepsilon a_n$, and $X \equiv X_1$, using the assumed condition (ii) of Theorem 1 to guarantee (18), and getting (19) and (20) (with an appropriately chosen constant) from (2) and (35) (which hold for all $n \geq 2$ by assumption), respectively. The Lemma then yields (33) since $r \geq \theta > 1$, and the proof is complete. $\square$

Proof of Theorem 1. Let $\alpha_1, \alpha_2, \dots$ be a sequence of independent Bernoulli random variables with $P(\alpha_n = 1) = P(\alpha_n = -1) = \frac{1}{2}$, and

with the sequence independent of $\{X_n\}_{n=1}^\infty$. Let $X'_n = \alpha_n X_n$, and put $S'_n = X'_1 + \cdots + X'_n$. Note that the primed variables are symmetric.

Suppose first that (4) holds and $\{\tau_n\}$ satisfies Condition A. By Lemma 2 we have (i) and (ii) holding. It remains to show that (iii) holds. But, if (4) holds, it likewise holds with S'_n in place of S_n , as can be seen by conditioning on the $\{\alpha_k\}_{k=1}^\infty$ and using [15, Corollary 5]. But then by the implication (a) $\Rightarrow$ (d) of Lemma 7 applied to the primed variables (which are symmetric) it follows that (iii) holds with X'_1 in place of X_1 . But (iii) holding for X'_1 is equivalent to it holding for X_1 , and so (iii) follows.

Conversely, suppose (i), (ii) and (iii) hold. Evidently (ii) and (iii) will also hold with X'_1 in place of X_1 . Hence by the implication (d) $\Rightarrow$ (a) of Lemma 7 as applied to the primed variables we have

$$(36) \quad \sum_{n=1}^{\infty} \tau_n P(|S'_n| \geq \varepsilon a_n) < \infty, \quad \forall \varepsilon > 0.$$

Now, for any random variable Y , let $Y^s = Y - \tilde{Y}$ be the symmetrization of Y , where $\tilde{Y}$ is an independent copy of Y . We shall choose symmetrizations in such a way that $X_1^s, X_2^s, \dots$ are independent and $S_n^s = X_1^s + \cdots + X_n^s$ for all n . Now, since

$$\begin{aligned} P(|X_k^s| \geq \lambda) &\leq P(|X_k| \geq \lambda/2) + P(|\tilde{X}_k| \geq \lambda/2) \\ &= 2P(|2X_k| \geq \lambda) = 2P(|2X'_k| \geq \lambda), \end{aligned}$$

for all $\lambda \geq 0$, it follows from [20, Theorem 1] that there is an absolute constant $c > 0$ such that

$$P(|S_n^s| \geq \lambda) \leq cP(|2S'_n| \geq \lambda/c)$$

for all $\lambda \geq 0$. By (36) it then follows that

$$\sum_{n=1}^{\infty} \tau_n P(|S_n^s| \geq \varepsilon a_n) < \infty, \quad \forall \varepsilon > 0.$$

By standard symmetrization inequalities [14, §17.1.A] it follows that

$$(37) \quad \sum_{n=1}^{\infty} \tau_n P(|S_n - \mu_n| \geq \varepsilon a_n) < \infty, \quad \forall \varepsilon > 0,$$

where μ_n is the median of S_n occurring in (i). Now, for any $\varepsilon > 0$ we have

$$(38) \quad \begin{aligned} \sum_{n=1}^{\infty} \tau_n P(|S_n| \geq \varepsilon a_n) \\ \leq \sum_{n=1}^{\infty} \tau_n P(|S_n - \mu_n| \geq \varepsilon a_n/2) + \sum_{n=1}^{\infty} \tau_n \cdot 1_{\{|\mu_n| > \varepsilon a_n/2\}}. \end{aligned}$$

By condition (i), the second sum converges, and by (37), so does the first, and hence (4) follows. $\square$

Proof of Theorem 2. Assume condition (ii) of Theorem 1 holds. Fix $\varepsilon > 0$. Changing a finite number of values of τ_n and a_n and using (5) and (6) will let us assume that if we let $b_n = \varepsilon a_n$ and $\nu = 2$, then conditions (19) and (20) of Lemma 5 will be verified for an appropriate constant. Applying that Lemma and using condition (ii) of Theorem 1

shows that:

$$(39) \quad \sum_{n=1}^{\infty} \tau_n n a_n^{-2} T_{\varepsilon, n} < \infty,$$

where $T_{\varepsilon, n}$ is as in Theorem 1(iii). Now, using the elementary inequality $e^{-1/x} \leq c_{\theta} x^{\theta}$ which is valid for all $x \geq 0$ where c_{θ} is a constant depending only on $\theta > 0$, we see that (39) entails that

$$\sum_{n=1}^{\infty} \tau_n e^{-\varepsilon^2 a_n^2 / (n T_{\varepsilon, n})} < \infty,$$

and so (iii) is true. $\square$

Proof of Lemma 1. Let $a_n = (n \log n)^{1/2}$ for $n \geq 2$. Assume the conditions of the Lemma hold. Without loss of generality $E[X_1] = 0$. Let $Y_n = X_1 \cdot 1_{\{|X_1| \leq a_n\}}$. Observe that Condition (b) of Corollary 4 implies that

$$(40) \quad \sum_{n=2}^{\infty} P(|X_1| > a_n) < \infty.$$

We now need the fact that in order to prove that $S_n/a_n \rightarrow 0$ in probability, given the fact that the X_k are identically distributed and with mean zero, all we need to show is that (40) holds as well as that

$$(41) \quad \lim_{n \rightarrow \infty} \frac{n E[Y_n^2]}{a_n^2} \rightarrow 0$$

and

$$(42) \quad \lim_{n \rightarrow \infty} \frac{n E[Y_n]}{a_n} \rightarrow 0.$$

To see this fact, let $\xi_{n,k} = X_k/a_n$ and $m_{n,k} = 0$ in the setting of [5, p. 105] (there $m_{n,k}$ is supposed to be a median of $\xi_{n,k}$, but the proof

in the direction we need does not use this fact), and note that (40)–(42) together with the assumption that $E[X_1] = 0$ (which assumption implies that $E[X_1 \cdot 1_{\{|X_1| > 1\}}] = -E[Y_1]$) and the proof in [5, p. 106] yield the convergence in probability of $(S_n - nE[Y_n])/a_n$ to zero, which in turn implies the convergence in probability of S_n/a_n to zero by (42).

Since as already noted, Condition (b) of Corollary 4 implies (40), we need only prove (41) and (42). First, observe that for $n \geq 2$ we have:

$$(43) \quad \begin{aligned} \frac{nE[Y_n^2]}{a_n^2} &= (\log n)^{-1} E[X_1^2 \cdot 1_{\{|X_1| \leq a_n\}}] \\ &= E \left[\frac{X_1^2}{\log^+ X_1} \cdot \frac{1_{\{|X_1| \leq a_n\}} \cdot \log^+ X_1}{\log n} \right]. \end{aligned}$$

Let

$$W_n = \frac{X_1^2}{\log^+ X_1} \cdot \frac{1_{\{|X_1| \leq a_n\}} \cdot \log^+ X_1}{\log n}.$$

Observe that

$$0 \leq W_n \leq c \frac{X_1^2}{\log^+ X_1},$$

where $c = \sup_{n \geq 2} (\log^+ a_n) / (\log n) < \infty$. But $X_1^2 / \log^+ X_1$ is in L^1 , while $W_n \rightarrow 0$ pointwise as $n \rightarrow \infty$, so that by dominated convergence it follows that $E[W_n] \rightarrow 0$ as $n \rightarrow \infty$ and hence (41) follows from (43).

Now, to prove (42), let $U_n = X_1 - Y_n$. Observe that $E[U_n] = -E[Y_n]$ since $E[X_1] = 0$. Hence, we only need to show that $a_n^{-1} n E[U_n] \rightarrow 0$. Now, since $|U_n| \geq a_n$ whenever $U_n \neq 0$ and as $x^{-1} \log^+ x$ is decreasing in $x > 0$, we have (using the convention that $(0^2 / \log^+ |0|) \cdot (\log^+ |0|) / |0| =$

0):

$$\begin{aligned}
a_n^{-1}nE[|U_n|] &= a_n^{-1}nE[(U_n^2/\log^+ |U_n|) \cdot (\log^+ |U_n|)/|U_n|] \\
&\leq a_n^{-1}nE[(U_n^2/\log^+ |U_n|) \cdot (\log^+ a_n)/a_n] \\
&= (\log^+ a_n)na_n^{-2}E[(U_n^2/\log^+ U_n)] \rightarrow 0,
\end{aligned}$$

since $(\log^+ a_n)na_n^{-2}$ is bounded while $E[(U_n^2/\log^+ U_n)] \rightarrow 0$ by dominated convergence, as $U_n \rightarrow 0$ almost surely and $U_n^2/\log^+ U_n \leq X_1^2/\log^+ X_1$, while $X_1^2/\log^+ X_1 \in L^1$ by Condition (b) of Corollary 4. $\square$

ACKNOWLEDGMENTS

The author is most grateful for a number of interesting e-mail conversations on these topics with Professors Aurel Spătaru and Stephen Montgomery-Smith. In particular, the author is very grateful to Professor Montgomery-Smith for permission to use his Example 1, and to Professor Spătaru for a number of useful comments on earlier versions of this paper, including pointing out several errors. The author also expresses his gratitude for a very careful reading to two referees, and thanks for pointing out a number of errors in earlier drafts.

REFERENCES

1. L. E. Baum and M. Katz, Convergence rates in the law of large numbers, *Trans. Amer. Math. Soc.* **120** (1965), 108–123. [**MR** 33 #6679]

2. N. H. Bingham, C. M. Goldie, and J. L. Teugels, "Regular Variation," Encyclopedia of Mathematics and Its Applications, vol. 27, Cambridge University Press, Cambridge, 1987. [MR 88i:26004]
3. Paul Erdős, On a theorem of Hsu and Robbins, *Ann. Math. Statist.* **20** (1949), 286–291. [MR 11,375b]
4. ———, Remark on my paper "On a theorem of Hsu and Robbins", *Ann. Math. Statist.* **21** (1950), 138. [MR 11,40f]
5. B. V. Gnedenko and A. N. Kolmogorov, "Limit Distributions for Sums of Independent Random Variables," revised ed., Addison-Wesley, Reading, MA, 1968. [MR 38 #1722]
6. A. Gut, Complete convergence for arrays, *Periodica Math. Hungarica* **25** (1992), 51–75. [MR 93m:60059]
7. J. Hoffman-Jørgensen, Sums of independent Banach space valued random variables, *Studia Math.* **52** (1974), 159–186. [MR 50 #8626]
8. P. L. Hsu and H. Robbins, Complete convergence and the law of large numbers, *Proc. Nat. Acad. Sci. U.S.A.* **33** (1947), 25–31. [MR 8,470e]
9. T.-C. Hu, F. Moricz, and R. L. Taylor, Strong law of large numbers for arrays of rowwise independent random variables, *Acta Math. Acad. Sci. Hungar.* **54** (1989), 153–162. [MR 91f:60061]
10. T.-C. Hu, D. Szynal, and A. I. Volodin, *A note on complete convergence for arrays*, Statist. Probab. Letters **38** (1998), 27–31. [MR 99e:60084]
11. T.-C. Hu and A. I. Volodin, *Addendum to: "A note on complete convergence for arrays" by Hu, Volodin and D. Szynal*, Statist. Probab. Lett. **47** (2000), 209–211. [CMP 1 747 108]
12. O. I. Klesov, Convergence of series of probabilities of large deviations of sums of independent identically distributed random variables, *Ukrain's'kiĭ Mat. Zhurn.*

- 45** (1993), 770–784 (Russian), English translation in: *Ukrainian Math. J.* **45** (1993), 845–862. [MR 96e:60046]
13. Deli Li, M. Bhaskara Rao, Tiefeng Jiang, and Xiangchen Wang, Convergence and almost sure convergence of weighted sums of random variables, *J. Theoret. Probab.* **8** (1995), 49–76. [MR 95k:60060]
14. Michel Loève, “Probability Theory,” 3rd ed., Van Nostrand, New York, 1963. [MR 34 #3596]
15. S. J. Montgomery-Smith, Comparison of sums of independent identically distributed random vectors, *Probab. Math. Statist.* **14** (1993), 281–285. [MR 96b:60042]
16. S. V. Nagaev, Some limit theorems for large deviations, *Theory Probab. Appl.* **10** (1965), 214–235. [MR 32 #3106]
17. Alexander R. Pruss, On Spătaru’s extension of the Hsu-Robbins-Erdős law of large numbers, *J. Math. Anal. Appl.* **199** (1996), 558–578. [MR 97g:60061]
18. ———, Randomly sampled Riemann sums and complete convergence in the law of large numbers for a case without identical distribution, *Proc. Amer. Math. Soc.* **124** (1996), 919–929. [MR 96f:60053]
19. ———, Remarks on summability of series formed from deviation probabilities of sums of independent identically distributed random variables, *Ukrains’kii Mat. Zhurn.* **48** (1996), 569–572, reprinted in: *Ukrainian Math. J.* **48** (1996), 631–635. [MR 98g:60063]
20. ———, Comparisons between tail probabilities of sums of independent symmetric random variables, *Ann. Inst. Poincaré Probab. Statist.* **33** (1997), 651–671. [MR 98j:60053]

DEPARTMENT OF PHILOSOPHY, GEORGETOWN UNIVERSITY, WASHINGTON,
DC 20057, U.S.A.

E-mail address: ap85@georgetown.edu